

Theoretical Overview of Hadronic Three-body B Decays

Hai-Yang Cheng

Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China

Charmless 3-body decays of B mesons are studied in a simple model based on the factorization approach. We have identified two distinct sources of nonresonant contributions: a small contribution from the tree transition and a large source of the nonresonant signal in the matrix elements of scalar densities, e.g., $\langle K\bar{K}|\bar{s}s|0\rangle$, induced from the penguin transition. This explains the dominance of the nonresonant background in $B \rightarrow KKK$ decays, the sizable nonresonant fraction in $K^-\pi^+\pi^-$ and $\bar{K}^0\pi^+\pi^-$ modes and the smallness of nonresonant rates in $B \rightarrow \pi\pi\pi$ decays. The seemingly huge discrepancy between BaBar and Belle for the nonresonant contribution in the decay $B^- \rightarrow K^-\pi^+\pi^-$ is now relieved. We have computed the resonant and nonresonant contributions to charmless 3-body decays and determined the rates for the quasi-two-body decays $B \rightarrow VP$ and $B \rightarrow SP$. Time-dependent CP asymmetries $\sin 2\beta_{\text{eff}}$ and A_{CP} in $K^+K^-K_S$, $K_S K_S K_S$, $K_S \pi^+ \pi^-$ and $K_S \pi^0 \pi^0$ modes are estimated.

I. INTRODUCTION

Recently many three-body B decay modes, for example, $K^+\pi^+\pi^-$, $K^0\pi^+\pi^-$, $K^+\pi^-\pi^0$, $K^+K^+K^-$, $K^0K^+K^-$, $K_S K_S K^+$ and $\pi^+\pi^+\pi^-$, have been observed at B factories with branching ratios of order 10^{-5} . The Dalitz plot analysis of 3-body B decays provides a nice methodology for extracting information on the unitarity triangle in the standard model. The three-body meson decays are generally dominated by intermediate vector and scalar resonances, namely, they proceed via quasi-two-body decays containing a resonance state and a pseudoscalar meson. Indeed, most of the quasi-two B decays are extracted from the analysis of three-body B decays using the Dalitz plot technique. Three-body hadronic B decays involving a vector meson or charmed meson also have been observed at B factories. In this talk I'll focus on charmless 3-body B decays.

Experimentally, there are two striking features for 3-body hadronic B decays:

- (i) large nonresonant fractions in penguin-dominated modes

It is known that the nonresonant signal in charm decays is small, less than 10% [1]. In the past few years,

some of the charmless B to 3-body decay modes have been measured at B factories and studied using the Dalitz plot analysis. We see from Table I that the nonresonant fraction is about $\sim 90\%$ in $B \rightarrow KKK$ decays, $\sim 17 - 40\%$ in $B \rightarrow K\pi\pi$ decays (smaller in the $K\pi\pi^0$ decay), and $\sim 14\%$ in the $B \rightarrow \pi\pi\pi$ decay. Hence, the nonresonant 3-body decays play an essential role in penguin-dominated B decays. While this is a surprise in view of the rather small nonresonant contributions in 3-body charm decays, it is not entirely unexpected because the energy release scale in weak B decays is of order 5 GeV, whereas the major resonances lie in the energy region of 0.77 to 1.6 GeV. Consequently, it is likely that 3-body B decays may receive sizable nonresonant contributions. It is important to understand and identify the underlying mechanism for nonresonant decays.

Nonresonant amplitudes in charm decays are usually assumed to be uniform in phase space. However, this is no longer true in B decays due to the large energy release in weak B decays. While both BaBar and Belle have adopted the parametrization

$$A_{\text{NR}} = (c_{12}e^{i\phi_{12}}e^{-\alpha s_{12}^2} + c_{13}e^{i\phi_{13}}e^{-\alpha s_{13}^2} + c_{23}e^{i\phi_{23}}e^{-\alpha s_{23}^2})(1 + b_{\text{NR}}e^{i(\beta + \delta_{\text{NR}})}) \quad (1)$$

to describe the non-resonant $B \rightarrow KKK$ amplitudes, they differ in the analysis of the nonresonant component in $B \rightarrow K\pi\pi$ decays. Belle still employed the above exponential parametrization to analyze the nonresonant contribution, but BaBar used the LASS parametrization to describe the $K\pi$ S -wave and the nonresonant component by a single amplitude suggested by the LASS collaboration to describe the scalar amplitude in elastic $K\pi$ scattering. Since the BaBar and Belle definitions of the $K_0^*(1430)$ and nonresonant differ, the branching fractions and phases are not directly comparable. We will come this point more in Sec. III.

Experimentally, it is hard to measure the direct 3-body decays as the interference between nonresonant and quasi-two-body amplitudes makes it difficult to

TABLE I: Fractions (%) of nonresonant contributions to various charmless three-body decays of B mesons [2]. It will be explained below about the BaBar measurement of the nonresonant component in the $B^- \rightarrow K^-\pi^+\pi^-$ decay.

Decay	BaBar	Belle
$B^- \rightarrow K^+K^-K^-$	141 ± 18	74.8 ± 3.6
$\bar{B}^0 \rightarrow K^+K^-\bar{K}^0$	112 ± 15	
$\bar{B}^0 \rightarrow \bar{K}^0\pi^+\pi^-$		41.9 ± 5.5
$B^- \rightarrow K^-\pi^+\pi^-$	$17.1^{+12.5}_{-2.5}$	34.0 ± 2.9
$\bar{B}^0 \rightarrow K^+\pi^-\pi^0$		$15.6 \pm 7.7 < 25.7$
$B^- \rightarrow \pi^+\pi^-\pi^-$	13.6 ± 6.1	

disentangle these two distinct contributions and extract the nonresonant one.

(ii) New broad scalar resonances $f_X(1550)$ and $f_X(1300)$

A broad scalar resonance $f_X(1500)$ (or $X_0(1550)$ denoted by BaBar) has been seen in $B \rightarrow K^+ K^+ K^-$, $K^+ K^- K_S$ and $K^+ K^- \pi^+$ decays at energies around 1.5 GeV. However, it cannot be identified with the well known scalar meson $f_0(1500)$. This is because $f_0(1500)$ decays into $\pi^+ \pi^-$ about five times more frequently than to $K^+ K^-$. Identification of $f_X(1500)$ with $f_0(1500)$ will imply that the $K^+ K^-$ peak at 1.5 GeV will be accompanied by a peak in $\pi^+ \pi^-$, which is not seen experimentally. Hence, the nature of $f_X(1500)$ is not clear.

Moreover, there exists a production puzzle for $f_X(1500)$. Both BaBar and Belle have seen a large fraction from $f_X(1500)$ in the decay $B^- \rightarrow K^+ K^- K^-$: $(121 \pm 19 \pm 6)\%$ by BaBar [3] and $(63.4 \pm 6.9)\%$ by Belle [4], whereas it is only about 4% seen by BaBar in $B^0 \rightarrow K^+ K^- K_S$ [5]. The puzzle is that why $f_X(1500)$ behaves so dramatically different in charged and neutral B decays to 3 kaons. It is not clear whether the large production of $f_X(1500)$ is a genuine effect or just a statistical fluctuation. Anyway, this issue should be clarified soon. Notice that Belle actually found two solutions for the fraction of $f_X(1500) K^-$ in $B^- \rightarrow K^+ K^+ K^-$ [4]: $(63.4 \pm 6.9)\%$ and $(8.21 \pm 1.94)\%$. The first solution is preferred by Belle. It is probably worth of re-examining the small solution.

II. THREE-BODY B DECAYS

In analog to two-body decays of heavy mesons which can be analyzed using the model-independent quark diagrammatic approach, three-body decays of the heavy mesons can be expressed in terms of some quark-graph amplitudes [6, 7] (see Fig. 1): $\mathcal{T}_1$ and $\mathcal{T}_2$, the color-allowed external W -emission tree diagrams; $\mathcal{C}_1$ and $\mathcal{C}_2$, the color-suppressed internal W -emission diagrams; $\mathcal{E}$, the W -exchange diagram; $\mathcal{A}$, the W -annihilation diagram; $\mathcal{P}_1$ and $\mathcal{P}_2$, the penguin diagrams, and $\mathcal{P}_a$, the penguin-induced annihilation diagram. The quark-graph amplitudes of various 3-body B decays $B \rightarrow \pi h^+ h^-$ and $B \rightarrow K h^+ h^-$ are summarized in Table I of [8]. As mentioned in [6], the use of the quark-diagram amplitudes for three-body decays are in general momentum dependent. This means that unless its momentum dependence is known, the quark-diagram amplitudes of direct 3-body decays cannot be extracted from experiment without making further assumptions. Moreover, the momentum dependence of each quark-diagram amplitude varies from channel to channel.

We take the decay $\bar{B}^0 \rightarrow K^+ K^- \bar{K}^0$ as an illustra-




FIG. 1: Quark diagrams for the three-body decays of heavy mesons, where Q denotes a heavy quark.

tion. Under the factorization approach, its decay amplitude consists of three distinct factorizable terms: (i) the transition process induced by $b \rightarrow s$ penguins, $\langle \bar{B}^0 \rightarrow K^+ \bar{K}^0 \rangle \times \langle 0 \rightarrow K^- \rangle$, (ii) the current-induced process through the tree $b \rightarrow u$ transition, $\langle \bar{B}^0 \rightarrow \bar{K}^0 \rangle \times \langle 0 \rightarrow K^+ K^- \rangle$, and (iii) the annihilation process $\langle \bar{B}^0 \rightarrow 0 \rangle \times \langle 0 \rightarrow K^+ K^- \bar{K}^0 \rangle$, where $\langle A \rightarrow B \rangle$ denotes a $A \rightarrow B$ transition matrix element.

A. Nonresonant background

For the transition process, the general expression of the nonresonant contribution has the form

$$\begin{aligned} & \langle K^-(p_3) | (\bar{s}u)_{V-A} | 0 \rangle \langle \bar{K}^0(p_1) K^+(p_2) | (\bar{u}b)_{V-A} | \bar{B}^0 \rangle^{NR} \\ &= -\frac{f_K}{2} [2m_3^2 r + (m_B^2 - s_{12} - m_3^2) \omega_+ \\ & \quad + (s_{23} - s_{13}) \omega_-], \end{aligned} \quad (2)$$

where $(\bar{q}_1 q_2)_{V-A} \equiv \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$. In principle, one can apply heavy meson chiral perturbation theory (HMChPT) to evaluate the form factors r , ω_+ and ω_- (for previous studies, see [9]). However, this will lead to too large decay rates in disagreement with experiment [10]. A direct calculation indicates that the branching ratio of $\bar{B}^0 \rightarrow K^+ K^- \bar{K}^0$ arising from the transition process alone is already at the level of 77×10^{-6} which exceeds the measured total branching ratio [2] of 25×10^{-6} . The issue has to do with the applicability of HMChPT. In order to apply this approach, two of the final-state pseudoscalars (K^+ and $\bar{K}^0$ in this example) have to be soft. The momentum of the soft pseudoscalar should be smaller than the chiral symmetry breaking scale Λ_χ of order $0.83 - 1.0$ GeV. For 3-body charmless B decays, the available phase space where chiral perturbation theory is applicable is only a small fraction of the whole Dalitz plot.

Therefore, it is not justified to apply chiral and heavy quark symmetries to a certain kinematic region and then generalize it to the region beyond its validity. If the soft meson result is assumed to be the same in the whole Dalitz plot, the decay rate will be greatly overestimated.

Recently we have proposed to parametrize the $b \rightarrow u$ transition-induced nonresonant amplitude given by Eq. (2) as [11]

$$A_{NR} = A_{NR}^{\text{HMChPT}} e^{-\alpha_{NR} p_B \cdot (p_1 + p_2)} e^{i\phi_{12}}, \quad (3)$$

so that the HMChPT results are recovered in the chiral limit $p_1, p_2 \rightarrow 0$. That is, the nonresonant amplitude in the soft meson region is described by HMChPT, but its energy dependence beyond the chiral limit is governed by the exponential term $e^{-\alpha_{NR} p_B \cdot (p_1 + p_2)}$. The unknown parameter α_{NR} can be determined from the data of the tree-dominated decay $B^- \rightarrow \pi^+ \pi^- \pi^-$.

In addition to the $b \rightarrow u$ tree transition, we need to consider the nonresonant contributions to the $b \rightarrow s$ penguin amplitude

$$\begin{aligned} A_1 &= \langle \bar{K}^0 | (\bar{s}b)_{V-A} | \bar{B}^0 \rangle \langle K^+ K^- | (\bar{u}u)_{V-A} | 0 \rangle, \\ A_2 &= \langle \bar{K}^0 | \bar{s}b | \bar{B}^0 \rangle \langle K^+ K^- | \bar{s}s | 0 \rangle. \end{aligned} \quad (4)$$

The 2-kaon creation matrix elements can be expressed in terms of time-like kaon current form factors as

$$\begin{aligned} \langle K^+(p_{K^+}) K^-(p_{K^-}) | \bar{q} \gamma_\mu q | 0 \rangle &= (p_{K^+} - p_{K^-})_\mu F_q^{K^+ K^-}, \\ \langle K^0(p_{K^0}) \bar{K}^0(p_{\bar{K}^0}) | \bar{q} \gamma_\mu q | 0 \rangle &= (p_{K^0} - p_{\bar{K}^0})_\mu F_q^{K^0 \bar{K}^0} \end{aligned} \quad (5)$$

The weak vector form factors $F_q^{K^+ K^-}$ and $F_q^{K^0 \bar{K}^0}$ can be related to the kaon electromagnetic (e.m.) form factors $F_{em}^{K^+ K^-}$ and $F_{em}^{K^0 \bar{K}^0}$ for the charged and neutral kaons, respectively. Phenomenologically, the e.m. form factors receive resonant and nonresonant contributions

$$\begin{aligned} F_{em}^{K^+ K^-} &= F_\rho + F_\omega + F_\phi + F_{NR}, \\ F_{em}^{K^0 \bar{K}^0} &= -F_\rho + F_\omega + F_\phi + F'_{NR}. \end{aligned} \quad (6)$$

The resonant and nonresonant terms in Eq. (6) can be determined from a fit to the kaon e.m. data. The non-resonant contribution to the matrix element $\langle K^+ K^- | \bar{s}s | 0 \rangle$ is given by

$$\begin{aligned} \langle K^+(p_2) K^-(p_3) | \bar{s}s | 0 \rangle^{NR} &\equiv f_s^{K^+ K^-}(s_{23}) \\ &= \frac{v}{3} (3F_{NR} + 2F'_{NR}) + \sigma_{NR} e^{-\alpha s_{23}}. \end{aligned} \quad (7)$$

The nonresonant σ_{NR} term is introduced for the following reason. Although the nonresonant contributions to f_s^{KK} and F_s^{KK} are related through the equation of motion, the resonant ones are different and not related *a priori*. As stressed in [12], to apply the equation of motion, the form factors should be

away from the resonant region. In the presence of the resonances, we thus need to introduce a nonresonant σ_{NR} term which can be constrained by the measured $\bar{B}^0 \rightarrow K_S K_S K_S$ rate and the $K^+ K^-$ mass spectrum [11].

B. Resonant contributions

Vector meson and scalar resonances contribute to the two-body matrix elements $\langle P_1 P_2 | V_\mu | 0 \rangle$ and $\langle P_1 P_2 | S | 0 \rangle$, respectively. They can also contribute to the three-body matrix element $\langle P_1 P_2 | V_\mu - A_\mu | B \rangle$. Resonant effects are described in terms of the usual Breit-Wigner formalism. More precisely,

$$\begin{aligned} \langle K^+ K^- | \bar{q} \gamma_\mu q | 0 \rangle^R &= \sum_i \langle K^+ K^- | V_i \rangle \\ &\times \frac{1}{m_{V_i}^2 - s - im_{V_i} \Gamma_{V_i}} \langle V_i | \bar{q} \gamma_\mu q | 0 \rangle, \\ \langle K^+ K^- | \bar{s}s | 0 \rangle^R &= \sum_i \langle K^+ K^- | S_i \rangle \\ &\times \frac{1}{m_{S_i}^2 - s - im_{S_i} \Gamma_{S_i}} \langle S_i | \bar{s}s | 0 \rangle, \end{aligned} \quad (8)$$

where $V_i = \phi, \rho, \omega, \dots$ and $S_i = f_0(980), f_0(1370), f_0(1500), \dots$. In this manner we are able to figure out the relevant resonances which contribute to the 3-body decays of interest and compute the rates of $B \rightarrow VP$ and $B \rightarrow SP$.

III. PENGUIN-DOMINATED $B \rightarrow KKK$ AND $B \rightarrow K\pi\pi$ DECAYS

As mentioned in the previous section, we employ the decays $\bar{B}^0 \rightarrow K^+ K^- \bar{K}^0$ and $K_S K_S K_S$ to fix the nonresonant parameter σ_{NR} to be

$$\sigma_{NR} = e^{i\pi/4} (3.36_{-0.96}^{+1.12}) \text{ GeV}. \quad (9)$$

It turns out that the nonresonant contribution arises dominantly from the transition process (88%) via the scalar-density-induced vacuum to $K\bar{K}$ transition, namely, $\langle K^+ K^- | \bar{s}s | 0 \rangle$, and slightly from the current-induced process (3%). Physically, this is because the decay $B \rightarrow KKK$ is dominated by the $b \rightarrow s$ penguin transition. The nonresonant background in $B \rightarrow KK$ transition does not suffice to account for the experimental observation that the penguin-dominated decay $B \rightarrow KKK$ is dominated by the nonresonant contributions. This implies that the two-body matrix element e.g. $\langle K\bar{K} | \bar{s}s | 0 \rangle$ induced by the scalar density should have a large nonresonant component.

We have considered other $B \rightarrow KKK$ decays such as $B^- \rightarrow K^+ K^- K^-$ and $B^- \rightarrow K^- K_S K_S$ and

TABLE II: Branching ratios (in units of 10^{-6}) of resonant and nonresonant (NR) contributions to $B^- \rightarrow K^- \pi^+ \pi^-$. Theoretical errors correspond to the uncertainties in (i) α_{NR} , (ii) m_s , F_0^{BK} and σ_{NR} , and (iii) $\gamma = (59 \pm 7)^\circ$. For the BaBar results, the branching fraction of $\bar{K}_0^{*0}(1430)\pi^-$ comes only from the Breit-Wigner component of the LASS parametrization, while the nonresonant contribution includes both the nonresonant part of the LASS shape and the phase-space nonresonant piece.

Decay mode	BaBar [13]	Belle [14]	Theory [11]
$\bar{K}^{*0}\pi^-$	$7.2 \pm 0.4 \pm 0.7^{+0.3}_{-0.5}$	$6.45 \pm 0.43 \pm 0.48^{+0.25}_{-0.35}$	$3.0^{+0.0+0.8+0.0}_{-0.0-0.7-0.0}$
$\bar{K}_0^{*0}(1430)\pi^-$	$32.0 \pm 1.2 \pm 2.7^{+9.1}_{-1.4} \pm 5.2$	$32.0 \pm 1.0 \pm 2.4^{+1.1}_{-1.9}$	$10.5^{+0.0+3.2+0.0}_{-0.0-2.7-0.1}$
$\rho^0 K^-$	$3.56 \pm 0.45 \pm 0.43^{+0.38}_{-0.15}$	$3.89 \pm 0.47 \pm 0.29^{+0.32}_{-0.29}$	$1.3^{+0.0+1.9+0.1}_{-0.0-0.7-0.1}$
$f_0(980)K^-$	$10.3 \pm 0.5 \pm 1.3^{+1.5}_{-0.4}$	$8.78 \pm 0.82 \pm 0.65^{+0.55}_{-1.64}$	$7.7^{+0.0+0.4+0.1}_{-0.0-0.8-0.1}$
NR	$9.3 \pm 1.0 \pm 1.2^{+6.7}_{-0.4} \pm 1.2$	$16.9 \pm 1.3 \pm 1.3^{+1.1}_{-0.9}$	$18.7^{+0.5+11.0+0.2}_{-0.6-6.3-0.2}$
Total	$54.4 \pm 1.1 \pm 4.6$	$48.8 \pm 1.1 \pm 3.6$	$45.0^{+0.3+16.4+0.1}_{-0.4-10.5-0.1}$

TABLE III: Branching ratios (in units of 10^{-6}) of resonant and nonresonant (NR) contributions to $\bar{B}^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$.

Decay mode	Belle [16]	Theory [11]
$K^{*-}\pi^+$	$5.6 \pm 0.7 \pm 0.5^{+0.4}_{-0.3}$	$2.1^{+0.0+0.5+0.3}_{-0.0-0.5-0.3}$
$K_0^{*-}(1430)\pi^+$	$30.8 \pm 2.4 \pm 2.4^{+0.8}_{-3.0}$	$10.1^{+0.0+2.9+0.1}_{-0.0-2.5-0.2}$
$\rho^0 \bar{K}^0$	$6.1 \pm 1.0 \pm 0.5^{+1.0}_{-1.1}$	$2.0^{+0.0+1.9+0.1}_{-0.0-0.9-0.1}$
$f_0(980)\bar{K}^0$	$7.6 \pm 1.7 \pm 0.7^{+0.5}_{-0.7}$	$7.7^{+0.0+0.4+0.0}_{-0.0-0.7-0.0}$
NR	$19.9 \pm 2.5 \pm 1.6^{+0.7}_{-1.2}$	$15.6^{+0.1+8.3+0.0}_{-0.1-4.9-0.0}$
Total	$47.5 \pm 2.4 \pm 3.7$	$42.0^{+0.3+15.7+0.0}_{-0.2-10.8-0.0}$

found that they are also dominated by the nonresonant contributions. Our predicted branching ratio $\mathcal{B}(B^- \rightarrow K^+ K^- K^-)_{\text{NR}} = (25.3^{+4.9}_{-4.5}) \times 10^{-6}$ [11] is in good agreement with the Belle measurement of $(24.0^{+3.0}_{-6.2}) \times 10^{-6}$, but a factor of 2 smaller than the BaBar result of $(50 \pm 6 \pm 4) \times 10^{-6}$ [3].

The resonant and nonresonant contributions to the decay $B^- \rightarrow K^- \pi^+ \pi^-$ are shown in Table II. We see that the calculated $K^* \pi$ and ρK rates are smaller than the data by a factor of $2 \sim 3$. This seems to be a generic feature of the factorization approach such as QCD factorization where the predicted penguin-dominated VP rates are too small compared to experiment. We shall return back to this point later.

At first sight, it appears that the nonresonant branching ratio $(2.4 \pm 0.5 \pm 1.3^{+0.3}_{-0.8}) \times 10^{-6}$ in $B^- \rightarrow K^- \pi^+ \pi^-$ obtained by BaBar [13] is much smaller than the one $(16.9 \pm 1.3 \pm 1.3^{+1.1}_{-0.9}) \times 10^{-6}$ measured by Belle [14]. However as mentioned in the Introduction, since the BaBar and Belle definitions of the $K_0^{*0}(1430)$ and nonresonant differ, it does not make sense to compare the branching fractions and phases directly. While Belle [14] employed the exponential parametrization Eq. (1) to describe the nonresonant contribution, BaBar [13] used the LASS parametrization to describe the $K\pi$ S -wave and the nonresonant component by a single amplitude suggested by the LASS collaboration

$$\mathcal{M} = \frac{m_{K\pi}}{q \cot \delta_B - iq}$$

$$+ e^{2i\delta_B} \frac{m_0 \Gamma_0 \frac{m_0}{q_0}}{(m_0^2 - m_{K\pi}^2) - im_0 \Gamma_0 \frac{q}{m_{K\pi}} \frac{m_0}{q_0}}, \quad (10)$$

where $\cot \delta_B = \frac{1}{aq} + \frac{1}{2}rq$. Since the LASS parametrization is valid (experimentally confirmed) up to the $K\pi$ invariant mass of order 1.8 GeV, BaBar introduced a phase-space nonresonant component to describe an excess of signal events at higher $K\pi$ invariant mass. Hence, the BaBar definition for the $K_0^{*0}(1430)$ includes an effective range term to account for the low $K\pi$ S -wave while for the Belle parameterization, this component is absorbed into the nonresonant piece. To stress once again, the result $\mathcal{B}(B^- \rightarrow K^- \pi^+ \pi^-)_{\text{NR}} = (2.4 \pm 0.5 \pm 1.3^{+0.3}_{-0.8}) \times 10^{-6}$ cited by BaBar is solely due to the phase-space nonresonant piece.

From the above discussion, it is clear that part of the LASS shape is really nonresonant which has a substantial mixing with $K_0^{*0}(1430)$. In principle, this should be added to the phase-space nonresonant piece to get the total nonresonant contribution. Once this is done, it is possible that BaBar and Belle might agree with each other. Indeed, very recently BaBar have carried out this task [13]. By combining coherently the nonresonant part of the LASS parametrization and the phase-space nonresonant, BaBar found the total nonresonant branching fraction to be $(9.3 \pm 1.0 \pm 1.2^{+6.7}_{-0.4} \pm 1.2) \times 10^{-6}$ with the fit fraction being $(17.1 \pm 1.7 \pm 1.6^{+12.3}_{-0.8})\%$ [15]. We see from Table II that the BaBar result is now

TABLE IV: Branching ratios (in units of 10^{-6}) of resonant and nonresonant (NR) contributions to $\bar{B}^0 \rightarrow K^- \pi^+ \pi^0$. For the BaBar measurement, the resonance $K_0^*(1430)$ is replaced by the S -wave $K\pi$ state, namely, $(K\pi)_0^*$.

Decay mode	BaBar [18]	Belle [19]	Theory [11]
$K^{*-} \pi^+$	$4.2^{+0.9}_{-0.5} \pm 0.3$	$4.9^{+1.5+0.5+0.8}_{-1.5-0.3-0.3}$	$1.0^{+0.0+0.3+0.1}_{-0.0-0.3-0.1}$
$\bar{K}^{*0} \pi^0$	$2.4 \pm 0.5 \pm 0.3$	< 2.3	$1.0^{+0.0+0.3+0.2}_{-0.0-0.2-0.1}$
$K_0^{*-}(1430) \pi^+$	$9.4^{+1.1+1.4}_{-1.3-1.1} \pm 1.8$	$5.1 \pm 1.5^{+0.6}_{-0.7}$	$5.0^{+0.0+1.5+0.1}_{-0.0-1.3-0.1}$
$\bar{K}_0^{*0}(1430) \pi^0$	$8.7^{+1.1+1.8}_{-0.9-1.3} \pm 2.2$	$6.1^{+1.6+0.5}_{-1.5-0.6}$	$4.2^{+0.0+1.4+0.0}_{-0.0-1.2-0.0}$
$\rho^+ K^-$	$8.0^{+0.8}_{-1.3} \pm 0.6$	$15.1^{+3.4+1.4+2.0}_{-3.3-1.5-2.1}$	$2.5^{+0.0+3.6+0.2}_{-0.0-1.4-0.2}$
NR		$5.7^{+2.7+0.5}_{-2.5-0.4} < 9.4$	$9.6^{+0.3+6.6+0.0}_{-0.2-3.5-0.0}$
Total	$35.7^{+2.6}_{-1.5} \pm 2.2$	$36.6^{+4.2}_{-4.1} \pm 3.0$	$28.9^{+0.2+16.1+0.2}_{-0.2-9.4-0.2}$

TABLE V: Same as Table II except for $B^- \rightarrow \pi^+ \pi^- \pi^-$. The nonresonant background is used as an input to fix the parameter α_{NR} defined in Eq. (3).

Decay mode	BaBar [20]	Theory [11]
$\rho^0 \pi^-$	$8.8 \pm 1.0 \pm 0.6^{+0.1}_{-0.7}$	$7.7^{+0.0+1.7+0.3}_{-0.0-1.6-0.2}$
$f_0(980) \pi^-$	$1.2 \pm 0.6 \pm 0.1 \pm 0.4 < 3.0$	$0.39^{+0.00+0.01+0.03}_{-0.00-0.01-0.02}$
NR	$2.3 \pm 0.9 \pm 0.3 \pm 0.4 < 4.6$	input
Total	$16.2 \pm 1.2 \pm 0.9$	$12.0^{+1.1+2.0+0.4}_{-1.2-1.8-0.3}$

consistent with Belle within errors, though the agreement is not perfect as BaBar and Belle have different models for the nonresonant $K\pi$ mass. Likewise, the BaBar branching fraction $(24.5 \pm 0.9 \pm 2.1^{+7.0}_{-1.1}) \times 10^{-6}$ for $B^- \rightarrow K_0^*(1430) \pi^-$ cited in Table II of [13] includes an effective range nonresonant component. In order to compare with the Belle result determined from the Breit-Wigner parametrization, it would be more appropriate to consider the Breit-Wigner component only of the LASS parametrization. The result is $\mathcal{B}(B^- \rightarrow K_0^*(1430) \pi^-) = (32.0 \pm 1.2 \pm 2.7^{+9.1}_{-1.4} \pm 5.2) \times 10^{-6}$ which is now in good agreement with the Belle measurement (see Table II).

From Table II we see that our predicted nonresonant rates are consistent with the Belle and BaBar measurements within errors. The reason for the large nonresonant rates in the $K^- \pi^+ \pi^-$ mode is that under SU(3) flavor symmetry, we have the relation $\langle K\pi | \bar{s}q | 0 \rangle^{NR} = \langle K\bar{K} | \bar{s}s | 0 \rangle^{NR}$. Hence, the nonresonant rates in the $K^- \pi^+ \pi^-$ (Table II) and $\bar{K}^0 \pi^+ \pi^-$ (Table III) modes should be similar to that in $K^+ K^- \bar{K}^0$ or $K^+ K^- K^-$. Since the KKK channel receives resonant contributions only from ϕ and f_0 mesons, while K^*, K_0^*, ρ, f_0 resonances contribute to $K\pi\pi$ modes, this explains why the nonresonant fraction is of order 90% in the former and becomes of order 40% in the latter. It is interesting to notice that, based on a simple fragmentation model and SU(3) symmetry, Gronau and Rosner [17] also found a large nonresonant background in $K^- \pi^+ \pi^-$ and $\bar{K}^0 \pi^+ \pi^-$.

Recently, BaBar has reported a new Dalitz-plot analysis of the decay $\bar{B}^0 \rightarrow K^- \pi^+ \pi^0$ [18] (see Table

IV). Just as the $K^- \pi^+ \pi^-$ mode, the reported nonresonant branching fraction $\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+ \pi^0)_{\text{nr}} = (4.4 \pm 0.9 \pm 0.5) \times 10^{-6}$ by BaBar is only the phase-space part of nonresonant contributions. To get the total nonresonant rate, it is necessary to add the nonresonant component of the LASS parametrization to the phase-space piece. When this is done, it will be interesting to compare the measured nonresonant branching fraction with our prediction $\mathcal{B}(\bar{B}^0 \rightarrow K^- \pi^+ \pi^0)_{\text{nr}} = (9.6^{+0.3+6.6}_{-0.2-3.5}) \times 10^{-6}$. It should be stressed that the measured partial rates for $\bar{B}^0 \rightarrow (K\pi)_0^{*-} \pi^+$ and $(K\pi)_0^{*0} \pi^0$ by BaBar (see Table IV) include an effective range $K\pi$ nonresonant component. Hence, it is not pertinent to compare them directly with the respective Belle measurements.

IV. TREE-DOMINATED $B \rightarrow \pi\pi\pi, K\bar{K}\pi$ MODES

The $B \rightarrow \pi\pi\pi$ mode receives nonresonant contributions mostly from the $b \rightarrow u$ transition as the nonresonant contribution in the penguin matrix element $\langle \pi^+ \pi^- | \bar{d}d | 0 \rangle$ is suppressed by the smallness of penguin Wilson coefficients a_6 and a_8 . Hence, the measurement of the nonresonant contribution in this decay can be used to constrain the nonresonant parameter α_{NR} in Eq. (3).

Note that while $B^- \rightarrow \pi^+ \pi^- \pi^-$ is dominated by the ρ^0 pole (Table V), the decay $\bar{B}^0 \rightarrow \pi^+ \pi^- \pi^0$ receives $\rho^\pm$ and ρ^0 contributions. As a consequence, the $\pi^+ \pi^- \pi^0$ mode has a rate larger than $\pi^+ \pi^- \pi^-$ even

though the former involves a π^0 in the final state. We predict that $\mathcal{B}(B^0 \rightarrow \pi^+\pi^-\pi^0) \approx 26 \times 10^{-6}$ [11].

Among the 3-body decays we have studied, the decay $B^- \rightarrow K^+K^-\pi^-$ dominated by $b \rightarrow u$ tree transition and $b \rightarrow d$ penguin transition has the smallest branching ratio of order 4×10^{-6} . BaBar [21] has recently reported the observation of the decay $B^+ \rightarrow K^+K^-\pi^+$ with the branching ratio $(5.0 \pm 0.5 \pm 0.5) \times 10^{-6}$. Our prediction for this mode, $(4.0_{-0.6-0.5-0.3}^{+0.5+0.7+0.3}) \times 10^{-6}$, is in accordance with experiment.

V. QUASI-TWO-BODY B DECAYS

It is known that in the narrow width approximation, the 3-body decay rate obeys the factorization relation

$$\Gamma(B \rightarrow RP \rightarrow P_1P_2P) = \Gamma(B \rightarrow RP)\mathcal{B}(R \rightarrow P_1P_2), \quad (11)$$

with R being a vector meson or a scalar resonance. Using the experimental information on $\mathcal{B}(R \rightarrow h_2h_3)$ [1]

$$\begin{aligned} \mathcal{B}(K^{*0} \rightarrow K^+\pi^-) &= \mathcal{B}(K^{*+} \rightarrow K^0\pi^+) \\ &= 2\mathcal{B}(K^{*+} \rightarrow K^+\pi^0) = \frac{2}{3}, \\ \mathcal{B}(K_0^{*0}(1430) \rightarrow K^+\pi^-) &= 2\mathcal{B}(K_0^{*+}(1430) \rightarrow K^+\pi^0) \\ &= \frac{2}{3}(0.93 \pm 0.10), \\ \mathcal{B}(\phi \rightarrow K^+K^-) &= 0.492 \pm 0.006, \end{aligned} \quad (12)$$

we have extracted the branching ratios of $B \rightarrow VP$ and $B \rightarrow SP$. The results are summarized in Table VI. The predicted $\rho\pi$, $f_0(980)K$ and $f_0(980)\pi$ rates are in agreement with the data, while the calculated ϕK , $K^*\pi$, ρK and $K_0^*(1430)\pi$ are in general too small compared to experiment. The fact that this work and QCDF lead to too small rates for ϕK , $K^*\pi$, ρK and $K_0^*(1430)\pi$ may imply the importance of power corrections due to the non-vanishing ρ_A and ρ_H parameters arising from weak annihilation and hard spectator interactions, respectively, which are used to parametrize the endpoint divergences, or due to possible final-state rescattering effects from charm intermediate states [22]. However, this is beyond the scope of the present work.

VI. TIME-DEPENDENT CP ASYMMETRIES

The penguin-induced three-body decays $B^0 \rightarrow K^+K^-K_S$ and $K_S K_S K_S$ deserve special attention as the current measurements of the deviation of $\sin 2\beta_{\text{eff}}$ in KKK modes from $\sin 2\beta_{J/\psi K_S}$ may indicate New Physics in $b \rightarrow s$ penguin-induced modes. It is of

great importance to examine and estimate how much of the deviation of $\sin 2\beta_{\text{eff}}$ is allowed in the SM. Owing to the presence of color-allowed tree contributions in $B^0 \rightarrow K^+K^-K_S$, this mode is subject to a potentially significant tree pollution and the deviation of the mixing-induced CP asymmetry from that measured in $B \rightarrow J/\psi K_S$ could be as large as $\mathcal{O}(0.10)$. Since the tree amplitude is tied to the nonresonant background, it is very important to understand the nonresonant contributions in order to have a reliable estimate of $\sin 2\beta_{\text{eff}}$ in KKK modes.

The deviation of the mixing-induced CP asymmetry in $B^0 \rightarrow K^+K^-K_S$, $K_S K_S K_S$, $K_S \pi^+ \pi^-$ and $K_S \pi^0 \pi^0$ from that measured in $B \rightarrow \phi_{c\bar{c}} K_S$, i.e. $\sin 2\beta_{\phi_{c\bar{c}} K_S} = 0.681 \pm 0.025$ [2], namely, $\Delta \sin 2\beta_{\text{eff}} \equiv \sin 2\beta_{\text{eff}} - \sin 2\beta_{\phi_{c\bar{c}} K_S}$, is shown in Table VII. Our calculation indicates the deviation of the mixing-induced CP asymmetry in $\bar{B}^0 \rightarrow K^+K^-K_S$ from that measured in $\bar{B}^0 \rightarrow \phi_{c\bar{c}} K_S$ is very similar to that of the $K_S K_S K_S$ mode as the tree pollution effect in the former is somewhat washed out. Nevertheless, direct CP asymmetry of the former, being of order -4% , is more prominent than the latter.

VII. CONCLUSIONS

It is important to account for the large nonresonant amplitudes in the study of charmless 3-body baryonic B decays. We have identified two distinct sources of nonresonant contributions: a small contribution from the tree transition and a large source of the nonresonant signal in the matrix elements of scalar densities, e.g. $\langle K\bar{K}|\bar{s}s|0\rangle$, induced from the penguin transition. This explains the dominance of the nonresonant background in $B \rightarrow KKK$ decays, the sizable nonresonant fraction in $K^-\pi^+\pi^-$ and $\bar{K}^0\pi^+\pi^-$ modes and the smallness of nonresonant rates in $B \rightarrow \pi\pi\pi$ decays. The seemingly huge discrepancy between BaBar and Belle for the nonresonant contribution in the decay $B^- \rightarrow K^-\pi^+\pi^-$ is now relieved. Since penguin contributions to charm decays are GIM suppressed, hence nonresonant signals in D decays are always small.

We have computed the resonant and nonresonant contributions to charmless 3-body decays and determined the rates for the quasi-two-body decays $B \rightarrow VP$ and $B \rightarrow SP$. Time-dependent CP asymmetries $\sin 2\beta_{\text{eff}}$ and A_{CP} in $K^+K^-K_S$, $K_S K_S K_S$, $K_S \pi^+ \pi^-$ and $K_S \pi^0 \pi^0$ modes are estimated. Since we have a realistic model for resonant and nonresonant contributions, our estimation of $\sin 2\beta_{\text{eff}}$ for 3-body B decays should be more reliable and trustworthy.

TABLE VI: Branching ratios of quasi-two-body decays $B \rightarrow VP$ and $B \rightarrow SP$ obtained from the studies of three-body decays based on the factorization approach [11]. Theoretical uncertainties have been added in quadrature. QCD factorization predictions taken from [23] for VP modes and from [24] for SP channels are shown here for comparison. The assumption of $\mathcal{B}(f_0(980) \rightarrow \pi^+\pi^-) = 0.50$ has been made for the QCDF calculation. Experimental results are taken from [2]. Note that the BaBar results for $\bar{K}_0^{*0}(1430)\pi^0$ and $K_0^{*-}(1430)\pi^+$ are obtained by neglecting nonresonant contributions to $(K\pi)_0^*\pi$ [18] and hence may not be appropriate to compare with Belle directly.

Decay mode	BaBar	Belle	QCDF	Theory [11]
ϕK^0	$8.4^{+1.5}_{-1.3} \pm 0.5$	$9.0^{+2.2}_{-1.8} \pm 0.7$	$4.1^{+0.4+1.7+1.8+10.6}_{-0.4-1.6-1.9-3.0}$	$5.3^{+1.0}_{-0.9}$
ϕK^-	$8.4 \pm 0.7 \pm 0.7$	$9.60 \pm 0.92^{+1.05}_{-0.84}$	$4.5^{+0.5+1.8+1.9+11.8}_{-0.4-1.7-2.1-3.3}$	$5.9^{+1.1}_{-1.0}$
$\bar{K}^{*0}\pi^-$	$10.8 \pm 0.6^{+1.1}_{-1.3}$	$9.7 \pm 0.6^{+0.8}_{-0.9}$	$3.6^{+0.4+1.5+1.2+7.7}_{-0.3-1.4-1.2-2.3}$	$4.4^{+1.1}_{-1.0}$
$\bar{K}^{*0}\pi^0$	$3.6 \pm 0.7 \pm 0.4$	< 3.5	$0.7^{+0.1+0.5+0.3+2.6}_{-0.1-0.4-0.3-0.5}$	$1.5^{+0.5}_{-0.4}$
$K^{*-}\pi^+$	$11.7^{+1.3}_{-1.2}$	$8.4 \pm 1.1^{+0.9}_{-0.8}$	$3.3^{+1.4+1.3+0.8+6.2}_{-1.2-1.2-0.8-1.6}$	$3.1^{+0.9}_{-0.9}$
$K^{*-}\pi^0$	$6.9 \pm 2.0 \pm 1.3$		$3.3^{+1.1+1.0+0.6+4.4}_{-1.0-0.9-0.6-1.4}$	$2.2^{+0.6}_{-0.5}$
$K^{*0}K^-$	< 1.1		$0.30^{+0.11+0.12+0.09+0.57}_{-0.09-0.10-0.09-0.19}$	$0.35^{+0.06}_{-0.06}$
$\rho^0 K^-$	$3.56 \pm 0.45^{+0.57}_{-0.46}$	$3.89 \pm 0.47^{+0.43}_{-0.41}$	$2.6^{+0.9+3.1+0.8+4.3}_{-0.9-1.4-0.6-1.2}$	$1.3^{+1.9}_{-0.7}$
$\rho^0 \bar{K}^0$	$4.9 \pm 0.8 \pm 0.9$	$6.1 \pm 1.0 \pm 1.1$	$4.6^{+0.5+4.0+0.7+6.1}_{-0.5-2.1-0.7-2.1}$	$2.0^{+1.9}_{-0.9}$
$\rho^+ K^-$	$8.0^{+0.8}_{-1.3} \pm 0.6$	$15.1^{+3.4+2.4}_{-3.3-2.6}$	$7.4^{+1.8+7.1+1.2+10.7}_{-1.9-3.6-1.1-3.5}$	$2.5^{+3.6}_{-1.4}$
$\rho^- \bar{K}^0$	$8.0^{+1.4}_{-1.3} \pm 0.6$		$5.8^{+0.6+7.0+1.5+10.3}_{-0.6-3.3-1.3-3.2}$	$1.3^{+3.0}_{-0.9}$
$\rho^0 \pi^-$	$8.8 \pm 1.0^{+0.6}_{-0.9}$	$8.0^{+2.3}_{-2.0} \pm 0.7$	$11.9^{+6.3+3.6+2.5+1.3}_{-5.0-3.1-1.2-1.1}$	$7.7^{+1.7}_{-1.6}$
$\rho^- \pi^+$			$21.2^{+10.3+8.7+1.3+2.0}_{-8.4-7.2-2.3-1.6}$	$15.5^{+4.0}_{-3.5}$
$\rho^+ \pi^-$			$15.4^{+8.0+5.5+0.7+1.9}_{-6.4-4.7-1.3-1.3}$	$8.5^{+1.1}_{-1.0}$
$\rho^0 \pi^0$	$1.4 \pm 0.6 \pm 0.3$	$3.0 \pm 0.5 \pm 0.7$	$0.4^{+0.2+0.2+0.9+0.5}_{-0.2-0.1-0.3-0.3}$	$1.0^{+0.3}_{-0.2}$
$f_0(980)K^0; f_0 \rightarrow \pi^+\pi^-$	$5.5 \pm 0.7 \pm 0.6$	$7.6 \pm 1.7^{+0.8}_{-0.9}$	$6.7^{+0.1+2.1+2.3}_{-0.1-1.5-1.1}$	$7.7^{+0.4}_{-0.7}$
$f_0(980)K^-; f_0 \rightarrow \pi^+\pi^-$	$9.3 \pm 1.0^{+0.6}_{-0.9}$	$8.8 \pm 0.8^{+0.9}_{-1.8}$	$7.8^{+0.2+2.3+2.7}_{-0.2-1.6-1.2}$	$7.7^{+0.4}_{-0.8}$
$f_0(980)K^0; f_0 \rightarrow K^+K^-$	5.3 ± 2.2			$5.8^{+0.1}_{-0.5}$
$f_0(980)K^-; f_0 \rightarrow K^+K^-$	$6.5 \pm 2.5 \pm 1.6$	< 2.9		$7.0^{+0.4}_{-0.7}$
$f_0(980)\pi^-; f_0 \rightarrow \pi^+\pi^-$	< 3.0		$0.5^{+0.0+0.2+0.1}_{-0.0-0.1-0.0}$	$0.39^{+0.03}_{-0.02}$
$f_0(980)\pi^-; f_0 \rightarrow K^+K^-$				$0.50^{+0.06}_{-0.04}$
$f_0(980)\pi^0; f_0 \rightarrow \pi^+\pi^-$			$0.02^{+0.01+0.02+0.04}_{-0.01-0.00-0.01}$	$0.010^{+0.003}_{-0.002}$
$\bar{K}_0^{*0}(1430)\pi^-$	$32.0 \pm 1.2^{+10.8}_{-6.0}$	$51.6 \pm 1.7^{+7.0}_{-7.4}$	$11.0^{+10.3+7.5+49.9}_{-6.0-3.5-10.1}$	$16.9^{+5.2}_{-4.4}$
$\bar{K}_0^{*0}(1430)\pi^0$	$13.1^{+1.6+2.7}_{-1.5-1.9} \pm 3.6$	$9.8 \pm 2.5 \pm 0.9$	$6.4^{+5.4+2.2+26.1}_{-3.3-2.1-5.7}$	$6.8^{+2.3}_{-1.9}$
$K_0^{*-}(1430)\pi^+$	$28.2^{+3.3+4.3}_{-4.1-3.3} \pm 5.2$	$49.7 \pm 3.8^{+4.0}_{-6.1}$	$11.3^{+9.4+3.7+45.8}_{-5.8-3.7-9.9}$	$16.2^{+4.7}_{-4.0}$
$K_0^{*-}(1430)\pi^0$			$5.3^{+4.7+1.6+22.3}_{-2.8-1.7-4.7}$	$8.9^{+2.6}_{-2.2}$
$K_0^{*0}(1430)K^-$	< 2.2			$1.3^{+0.3}_{-0.3}$

TABLE VII: Mixing-induced and direct CP asymmetries for various charmless 3-body B decays. Experimental results are taken from [2].

Decay	$\sin 2\beta_{\text{eff}}$	$\Delta \sin 2\beta_{\text{eff}}$	Expt	$A_f(\%)$	Expt
$K^+K^-K_S$	$0.728^{+0.001+0.002+0.009}_{-0.002-0.001-0.020}$	$0.041^{+0.028}_{-0.033}$	0.05 ± 0.11	$-4.63^{+1.35+0.53+0.40}_{-1.01-0.54-0.34}$	-7 ± 8
$K_S K_S K_S$	$0.719^{+0.000+0.000+0.008}_{-0.000-0.000-0.019}$	$0.039^{+0.027}_{-0.032}$	-0.10 ± 0.20	$0.69^{+0.01+0.01+0.05}_{-0.01-0.03-0.07}$	14 ± 15
$K_S \pi^0 \pi^0$	$0.729^{+0.000+0.001+0.009}_{-0.000-0.001-0.020}$	$0.049^{+0.027}_{-0.032}$	-1.20 ± 0.41	$0.28^{+0.09+0.07+0.02}_{-0.06-0.06-0.02}$	-18 ± 22
$K_S \pi^+ \pi^-$	$0.718^{+0.001+0.017+0.008}_{-0.001-0.007-0.018}$	$0.038^{+0.031}_{-0.032}$		$4.94^{+0.03+0.03+0.32}_{-0.02-0.05-0.40}$	

Acknowledgments

I'm grateful to Chun-Khiang Chua and Amarjit Soni for fruitful collaboration, to Tim Gershon and

Jim Smith for discussions and to Paoti Chang and Hsiang-nan Li for organizing this stimulating conference.

-
- [1] Particle Data Group, Y.M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
 - [2] Heavy Flavor Averaging Group, <http://www.slac.stanford.edu/xorg/hfag>.
 - [3] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **74**, 032003 (2006).
 - [4] A. Garmash *et al.* (Belle Collaboration), Phys. Rev. D **71**, 092003 (2005).
 - [5] B. Aubert *et al.* (BaBar Collaboration), hep-ex/0607112.
 - [6] L.L. Chau and H.Y. Cheng, Phys. Rev. D **41**, 1510 (1990).
 - [7] L.L. Chau and H.Y. Cheng, Phys. Rev. D **36**, 137 (1987); Phys. Lett. B **222**, 285 (1989).
 - [8] H.Y. Cheng and K.C. Yang, Phys. Rev. D **66**, 054015 (2002).
 - [9] S. Fajfer, R.J. Oakes, and T.N. Pham, Phys. Rev. D **60**, 054029 (1999); Phys. Lett. B **539**, 67 (2002); S. Fajfer, T.N. Pham, and A. Prapotnik, Phys. Rev. D **70**, 034033 (2004).
 - [10] H.Y. Cheng and K.C. Yang, Phys. Rev. D **66**, 054015 (2002).
 - [11] H.Y. Cheng, C.K. Chua, and A. Soni, Phys. Rev. D **76**, 094006 (2007).
 - [12] H.Y. Cheng, C.K. Chua, and A. Soni, Phys. Rev. D **72**, 094003 (2005).
 - [13] B. Aubert *et al.* (BaBar Collaboration), arXiv:0803.4451 [hep-ex].
 - [14] A. Garmash *et al.* (Belle Collaboration), Phys. Rev. Lett. **96**, 251803 (2006); Phys. Rev. D **75**, 012006 (2007).
 - [15] I would like to thank Tim Gershon for providing me this number.
 - [16] A. Garmash *et al.* (Belle Collaboration), Phys. Rev. D **75**, 012006 (2007).
 - [17] M. Gronau and J.L. Rosner, Phys. Rev. D **72**, 094031 (2005).
 - [18] B. Aubert *et al.* (BaBar Collaboration), arXiv:0711.4417v2 [hep-ex].
 - [19] P. Chang *et al.* (Belle Collaboration), Phys. Lett. B **599**, 148 (2004).
 - [20] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **72**, 052002 (2005).
 - [21] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **99**, 221801 (2007).
 - [22] H.Y. Cheng, C.K. Chua, and A. Soni, Phys. Rev. D **71**, 014030 (2005).
 - [23] M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
 - [24] H.Y. Cheng, C.K. Chua, and K.C. Yang, Phys. Rev. D **73**, 014017 (2006).